

Multiple Sclerosis | Neurology Prescription Form

PATIENT INFORMATION You may also fax demographics/face sheet			INSURANCE INFO. Please fax copy of ALL insurance ca	rds *fro	nt & back
Patient Name:DOB:			Primary Insurance:		
			Policy Number:Group Number:		
City, State, Zip:	Cell Phone	<u> </u>	Rx Bin: Rx PCN:		
			Secondary Insurance:		
	Weight:lbs		Policy Number:Group Number:		
	•				
Emergency Contact Pr	none:		Rx Bin: Rx PCN:		
PRESCRIBER INFO	DRMATION				
			I#: DEA#: TAX ID:		
			y, State, Zip:		
Office Contact:		Pho	one: Fax:		
DIAGNOSIS INFOR	MATION Please fax recent l	abs. clinical notes. etc. to he	Ip expedite the prior authorization process		
□ Multiple Sclerosis Primary ICD-10: Date of First Demyelinating Event: Type: □ Clinically Isolated Syndrome □ Relapsing-Remitting □ Secondary Progressive □				psing	
			omorbidities:		
	S:		Primary ICD-10: Secondary ICD-10: Reasons for Discontinuation:		
,			neasons for discontinuation.		
MEDICATIONS: You	may tape Prescriptions here property DOSE/STRENGTH	rior to faxing	SIG	QTY	REFILLS
□Avonex®	☐ 30 mcg Prefilled Syringe #4	☐ Inject 30 mcg IM once weekly	Viu	Q.I.	TILI ILLO
LAVOITEX	☐ 30 mcg Pen #4				
□Betaseron® □Extavia®		□ Dose Titration: • Week 1-2: Inject 0.0625 mg/0.25 ml subcutaneously QOD • Week 3-4: Inject 0.125 mg/0.50 ml subcutaneously QOD			
	□ 0.3 mg vial		75 ml subcutaneously QOD • Week 7+: Inject 0.25 mg/1 ml subcutaneously QOD		
	☐ Maintenance Dose: 0.25 mg /1 ml subcutaneously QOD☐ Other:				
□Copaxone® (DAW)	☐ 20 mg Prefilled Syringe	□ 20 mg SQ QD	t 40 haura apart an tha sama 2 days asah wasi		
	☐ 40 mg Prefilled Syringe	□ 40 mg SQ 3 times a week, at least 48 hours apart on the same 3 days each week			
□Dalfampridine®	□ 10 mg ER tablet	□ 10 mg orally twice daily, approximately 12 hours apart			
□Dimethyl Fumarate	☐ 120 mg DR capsule ☐ 240 mg DR capsule	☐ Initial: 120 mg orally twice a day, for 7 days ☐ Maintenance: 240 mg orally twice a day, starting at day 8			
□Gilenya®	□ 0.5 mg capsule	☐ Take 0.5mg po QD			
□Glatiramer		☐ 20 mg SQ QD			
Acetate	☐ 20 mg Prefilled Syringe☐ 40 mg Prefilled Syringe	☐ 40 mg SQ 3 times a week, at leas	st 48 hours apart on the same 3 days each week		
□WhisperJECT [™] (autoinjector)		Autoinjector - to be used with Giat	tiramer Acetate injection, prefilled, glass syringe		
□Glatopa®	□ 00 ··· D. (III. 10 ···	□ 20 mg SQ QD			
□GlatopaJECT™	20 mg Prefilled Syringe40 mg Prefilled Syringe	☐ 40 mg SQ 3 times a week, at leas☐ Autoinjector - to be used with Glat	st 48 hours apart on the same 3 days each week		
(autoinjector)		-			
□Kesimpta®	☐ 20 mg Sensoready® Pen	☐ Initial: 20 mg SQ once weekly for 3 doses (weeks 0, 1, and 2) ☐ Maintenance: 20 mg SQ once monthly, starting at week 4			
□Rebif®	☐ Titration Dook (9.9mog/22mog)		ree times a week weeks 1-2, 22 mcg subcutaneously three times a week weeks 3-4,		
(prefilled syringe)	☐ Titration Pack (8.8mcg/22mcg) ☐ 22 mcg	☐ Maintenance: Inject 22mcg (0.5r	e times a week weeks 5+ (48 hours apart) nl) SQ three times a week (48 hours apart)		
□Rebif Rebidose® (pen)	☐ 44 mcg	☐ Maintenance: Inject 44mcg (0.5n☐ Other:	nl) SQ three times a week (48 hours apart)		
□Tetrabenazine	☐ 12.5 mg tablet	☐ 12.5 mg initial dose once daily oral	lly in the morning, may increase to 12.5 mg twice a day after 1 week (max 100 mg/day)		
Lieuabeliazille	☐ 25 mg tablet	□ Other:			
	□ 0.23 mg capsule □ Starter Kit: Days 1-4: Take 0.23 mg capsu		: 0.23 mg capsule by mouth once daily • Days 5-7: Take 0.46 mg capsule by mouth once daily capsule by mouth once daily • Days 5-7: Take 0.46 mg capsule by mouth once daily		
□Zeposia®	☐ 0.46 mg capsule ☐ 0.92 mg capsule		Take 0.92 mg capsule by mouth once daily		
	U.32 mg capsule	Other:	ng capadio by illuditi office daily		
□Other					
DELIVERY INFORMATION	\ \				
Need by:		o: ☐ Patient's home ☐ MD Office/ C	Clinic Other:		
PRESCRIBER'S SIGNATU	JRE REQUIRED				
MD NP PA Signature:	·		□ DAW Date:		<u></u>

*Signature on this form also provides consent to contact this patient's insurance provider for this prescription on the provider's behalf.